

FIG. 1.




FIG. 2.




FIG. 3.




FIG. 4A.




FIG. 4B.




FIG. 5A.




FIG. 5B.




FIG. 5C.




FIG. 5D.




FIG. 5E.




FIG. 5F.




FIG. 6A.




FIG. 6B.




FIG. 7A.




FIG. 7B.




FIG. 8A.




FIG. 9.




FIG. 10A.




FIG. 10B.




FIG. 10C.






FIG. 10D.

FIG. 11.



1ST DIMENSION: ISOELECTRIC FOCUSING



2ND DIMENSION: SDS PAGE

FIG. 12.




FIG. 13.




FIG. 14.





FIG. 15.







FIG. 17.

Fig. 18







FIG. 20







FIG. 22



Sample	Protein	Area	Intensity
Disease	Int std	1139.044	3.340518
	marker	695.4804	0.580081
Drug treated	Int std	1383.883	3.780178
	marker	71.05119	0.143227


Fig. 22D

Normalized marker peak area (marker/internal std):

Control 0.0
Disease 0.61
Drug treated 0.05

+

FIG. 23
Retentate Map™ of Human Urine
SELDI ProteinChip™ Array



+

FIG. 24




FIG. 25A.




FIG. 25B.




FIG. 26A.




FIG. 26B.




FIG. 26C.



FIG. 26D.




FIG. 27A.





FIG. 27B.






FIG. 28A.




FIG. 28B.







FIG. 28C.



MOLECULAR WEIGHT




FIG. 30.




FIG. 31.




FIG. 32.

FIG. 33A.




FIG. 33B.

